

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 020 517 A1**

(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
19.07.2000 Bulletin 2000/29

(21) Application number: 98931048.7

(22) Date of filing: 10.07.1998

(51) Int. Cl.<sup>7</sup>: **C12N 15/12**, **C12P 21/02**,  
**C12N 5/08**, **C07K 14/47**,  
**C07K 16/18**

(86) International application number:  
PCT/JP98/03106

(87) International publication number:  
WO 99/02677 (21.01.1999 Gazette 1999/03)

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**

(30) Priority: 11.07.1997 JP 20222797

(71) Applicant:  
**CHUGAI SEIYAKU KABUSHIKI KAISHA**  
Tokyo, 115-8543 (JP)

(72) Inventors:  
• **KATO, Yukio**  
Hiroshima-shi, Hiroshima 732-0062 (JP)  
• **KAWAMOTO, Takeshi**  
Yoshiki-gun, Yamaguchi 754-1200 (JP)

(74) Representative: **HOFFMANN - EITLE**  
Patent- und Rechtsanwälte  
Arabellastrasse 4  
81925 München (DE)


(54) **GENE ORIGINATING IN HUMAN CHONDROCYTE**

(57) This invention provides a gene specifically expressed in differentiated human chondrocytes. By culturing the chondrocytes in the presence of dibutyryl cAMP, the chondrocytes are cultured in a differentiated state and any gene is searched for which has a distinction in expression between differentiated chondrocytes and dedifferentiated chondrocytes. This allows the gene specifically expressed in the former to be obtained.

**Fig.2B**



**Fig.2A**



EP 1 020 517 A1

## Description

## TECHNICAL FIELD

5 [0001] This invention relates to a gene expressed specifically in differentiated chondrocytes originating in human (or human chondrocytes), a protein encoded by the gene, an antibody capable of binding to the protein, a method for culturing human chondrocytes in a differentiated state, and human chondrocytes that have been cultured by the method.

## 10 BACKGROUND ART

[0002] Searching for genes expressed specifically in chondrocytes in a differentiated state and analysis of the properties of the chondrocytes are not only important in analyzing the mechanism of differentiation and degeneration of cartilage, but also are indispensable for developing gene therapy for osteoarthritis and rheumatoid arthritis.

15 [0003] However, any method for monolayer culturing human chondrocytes in a differentiated state has not yet been established, although culture systems for rabbit or chicken chondrocytes in a differentiated state have been developed. (Kato et al. Proc. Natl. Acad. Sci. USA 85, 9552-9556 (1988); Oakes et al. J. Embryol. Exp. Morphol. 38, 239-263 (1977).) It is recognized that human chondrocytes maintain their differentiated phenotype in agarose gel (Benya P.D. and Shaffer J.D., Cell 30, 215-224 (1982)), but that they easily lose the differentiated phenotype in the monolayer culture which facilitates handling of cells. Accordingly, it is difficult to search for genes that are expressed specifically in human chondrocytes in a differentiated state thereof; there has not been provided any cell culture system useful in analyzing the properties of the chondrocytes in a differentiated state thereof.

## DISCLOSURE OF INVENTION

25 [0004] A principle object of this invention is to establish a method of monolayer culture for human chondrocytes in a differentiated state and further to obtain a gene expressed specifically in the chondrocytes in a differentiated state thereof.

[0005] The present inventors found that chondrocytes could be cultured in a differentiated state by culturing the chondrocytes in the presence of a certain compound; and in addition, they searched for genes having a distinction in expression between differentiated chondrocytes and dedifferentiated chondrocytes and discovered those which were specifically expressed in the former. Thus, this invention has been accomplished.

[0006] Particularly, this invention provides a DNA encoding a protein defined in (a) or (b) as described below: the DNA may be referred to as "DNA of this (the) invention" hereinbelow.

- 35 (a) A protein comprising an amino acid sequence set forth in SEQ ID NO: 2.  
 (b) A protein comprising an amino acid sequence derived from the amino acid sequence set forth in SEQ ID NO: 2 by deletion or substitution of one or more amino acids therefrom, or by addition of one or more amino acids thereto and capable of binding to nucleotide sequence CANNTG and/or nucleotide sequence CACNAG upon formation of a dimer,  
 40 wherein the amino acid sequence of a part of said protein corresponding to an amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2 is provided with not less than 85% of homology to the amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2. Preferably, DNA of this invention is a DNA defined in the following (c) or (d):  
 45 (c) A DNA comprising a nucleotide sequence of from nucleotide no. 207 to nucleotide no. 1442 of the nucleotide sequence set forth in SEQ ID NO: 1 or a complementary nucleotide sequence thereto.  
 (d) A DNA capable of hybridizing to the DNA defined in (c) under stringent conditions.

[0007] This invention also provides a protein encoded by DNA of the invention, as well as an antibody capable of binding to the protein.

50 [0008] Further, the invention provides a method for culturing human chondrocytes, which comprises monolayer culturing the chondrocytes in the presence of a membrane-permeable cAMP analog in an amount sufficient to cause the chondrocytes to maintain a differentiated state thereof as cartilage: the method may be referred to as "the culturing method of this (the) invention" hereinbelow. Preferably, the membrane-permeable cAMP analog is dibutyl cAMP (which may be denoted "dbcAMP" hereinbelow).

55 [0009] Still further, the invention provides human chondrocytes that have been cultured by the culturing method of the invention and that possess the properties defined in the following (1)-(3):

- (1) Exhibit a spherical shape and are abundant in extracellular matrix;
- (2) Can be stained with toluidine blue satisfactorily; and
- (3) DNA of the invention is expressed therein.

- 5 [0010] The DNA of this invention is believed to encode a novel transcription factor of the basic helix-loop-helix type (bHLH), and is predicted to play an important role such as the regulation of expression of various genes in the differentiation of cartilage. Therefore, a DNA of this invention, a protein encoded by the DNA, and an antibody capable of binding to the protein are useful in the analysis of mechanism of the differentiation and degeneration of cartilage, as well as in the development of gene therapy for osteoarthritis and rheumatoid arthritis.
- 10 [0011] According to the culturing method of this invention, chondrocytes can be monolayer cultured in a good differentiated state, which makes it easy to search for genes having a distinction in expression between chondrocytes in a differentiated state thereof and chondrocytes in a dedifferentiated state thereof; namely, to search for genes expressed specifically in chondrocytes in a differentiated state thereof. Also, an analysis of the properties of chondrocytes in a differentiated state thereof will be facilitated.

#### 15 BREIF DESCRIPTION OF DRAWINGS

##### [0012]

- 20 Fig. 1 shows a comparison between DEC1 and other bHLH factors in the bHLH region.  
 Fig. 2A is a photomicrograph showing the morphology of human chondrocytes in monolayer culture (a-no addition of dbcAMP, b-addition of dbcAMP).  
 Fig. 2B is a photograph showing the morphology of human chondrocytes in monolayer culture (the morphology of an organism) that were stained with toluidine blue (a-no addition of dbcAMP, b-addition of dbcAMP).  
 25 Fig. 3 is a graph showing that DEC1 mRNA was induced in a fibroblast strain, MRC5, originating in the human lung after addition of dbcAMP.  
 Fig. 4 is a graph showing that DEC1 mRNA was induced in Hela cells originating in human uterine cancer after addition of dbcAMP.  
 Fig. 5 is a graph showing that EC1 mRNA was induced in rabbit chondrocytes after addition of PTH.  
 30 Fig. 6 is a graph showing that DEC1 mRNA was induced in a rabbit chondrocyte culture system after addition of dbcAMP.  
 Fig. 7 is a graph showing that DEC1 mRNA was induced in a renal cell strain after addition of dbcAMP.

#### BEST MODE FOR CARRYING OUT THE INVENTION

- 35 [0013] Embodiments of this invention will be explained hereinbelow.
- [0014] As will be later shown in the Examples, any genes expressed specifically in chondrocytes in a differentiated state were searched for; and as a result, the amino acid sequence set forth in SEQ ID NO: 2 has been revealed for the first time. As used in the present specification, the protein having this amino acid sequence may be referred to as  
 40 "DEC1."
- [0015] Through a homology search using a protein database, it was found that DEC1 has a basic helix-loop-helix (bHLH) region (amino acid nos. 51-108 in SEQ ID NO: 2). It is known that a bHLH protein forms a dimer and binds to E box (CANNTG).
- [0016] Particularly, in this bHLH region, rat HES1 (47.5%), rat HES2 (42.6%), rat HES3 (40.3%), rat HES5 (37.7%),  
 45 Drosophila Hairy (abbreviated as "hairy") (39.3%), and Enhancer of Split m7 (abbreviated as "E(spl)m7") (37.7%) showed high homology: numerical values in the parentheses represent homology levels. Fig. 1 shows corresponding bHLH regions of the respective proteins for comparison. The conserved residues are enclosed by the frames.
- [0017] The HES family, hairy, and E(spl)m7 function as negative regulators which repress transcription by binding to N box (CACNAG). (Sasai et al. Genes & Dev. 6, 2620-2634 (1992); Ishibashi et al. Eur. J. Biochem. 215, 645-652 (1993); Akazawa et al. J. Biol. Chem. 267, 21879-21885 (1992); Ohsako et al. Genes & Dev. 8, 2743-2755 (1994); Dawson et al. Mol. Cell. Biol. 15, 6923-6931 (1995); Jan et al. Cell 75, 827-830 (1993).) They also have the Trp-Arg-Pro-Trp (WRPW) domain (SEQ ID NO: 3) at their C-terminus: the domain is believed to lead to suppression of a certain activator by a repressor (Dawson et al. Mol. Cell. Biol. 15, 6923-6931 (1995)). Although DEC1 resembles a bHLH factor, it does not have this WRPW domain. Thus, DEC1 is believed to be a novel transcription factor that modulates chondrogenesis.  
 55 In view of the foregoing, it is thought that DEC1 is able to bind to E box, as well as to N box.
- [0018] Accordingly, the DNA of this invention encompasses: a DNA that encodes a protein comprising the amino acid sequence set forth in SEQ ID NO: 2; and in addition, a protein comprising an amino acid sequence derived from the amino acid sequence set forth in SEQ ID NO: 2 by deletion or substitution of one or more amino acids therefrom,

or by addition of one or more amino acids thereto and capable of binding to nucleotide sequence CANNTG and/or nucleotide sequence CACNAG (preferably, nucleotide sequence CANNTG) upon formation of a dimer,

wherein the amino acid sequence of a part of said protein corresponding to an amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2 is provided with not less than 85% (preferably, not less than 90%) of homology to the amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2. Here, "N" in the above-described sequence represents A, G, C, or T. For an example of nucleotide sequence CANNTG, mentioned are CACGTG, CAGGTG, CAGTTG, and CACCTG. The expression, "encoding a protein," means that either one of complementary double strands has a nucleotide sequence encoding the protein when the DNA is double-stranded.

[0019] Substitution, deletion, or insertion of (an) amino acid residue(s) can be generated by introducing into a nucleotide sequence, variation such as the substitution, deletion, or insertion of nucleotide, according to a known method (e.g., site-specific mutation). Methods for determining the activity which allow the binding to nucleotide sequence CANNTG or CACNAG when the dimer is formed are known: for example, see Ohsako et al. *Genes & Dev.* 8, 2743-2755 (1994). One skilled in the art can readily select such substitution, deletion, or insertion of one or more amino acid residues that would not substantially impair this activity.

[0020] Concrete examples of DNA of this invention encompass DNAs defined in the following (c) and (d):

(c) A DNA comprising a nucleotide sequence of from nucleotide no. 207 to nucleotide no. 1442 of the nucleotide sequence set forth in SEQ ID NO: 1 or a complementary nucleotide sequence thereto.

(d) A DNA capable of hybridization to the DNA defined in (c) under stringent conditions.

[0021] Here, the "stringent conditions" means the conditions under which a so-called specific hybrid is formed, but any non-specific hybrid is not formed. These conditions may be difficult to be accurately expressed as numerical values; and for example, mentioned is a temperature in the range of from  $T_m$  to  $T_m$  minus 20 ° C where the  $T_m$  is the one for a perfectly matched hybrid, such as that between highly homologous nucleic acids, or alternatively the conditions under which DNAs having homology of not less than 80% hybridize to each other, but nucleic acids having homology lower than that do not hybridize to each other.

[0022] The DNA of this invention is preferably one that encodes the amino acid set forth in SEQ ID NO: 2, and is more preferably the DNA of (c) described above.

[0023] As will be later shown in the Examples, one of the nucleotide sequences for DNA of the invention has been determined; therefore, it is possible to synthesize the DNA based on this sequence. The DNA can also be obtained from chromosomal DNAs by PCR or hybridization using oligonucleotides or probes that have been prepared based on this very nucleotide sequence. Alternatively, the DNA can further be obtained either by carrying out RT-PCR with cartilage mRNA or by screening a cDNA library, such as cartilage, with polynucleotides having a nucleotide sequence that encodes the whole or a part of DEC1 as probes.

[0024] The protein encoded by DNA of this invention is a protein defined in (e) or (f) described below: the protein may be referred to as "the protein(s) of this (the) invention" hereinbelow.

(e) A protein comprising an amino acid sequence set forth in SEQ ID NO: 2.

(f) A protein comprising an amino acid sequence derived from the amino acid sequence set forth in SEQ ID NO: 2 by deletion or substitution of one or more amino acids therefrom, or by addition of one or more amino acids thereto and capable of binding to nucleotide sequence CANNTG and/or nucleotide sequence CACNAG upon formation of a dimer,

wherein the amino acid sequence of a part of said protein corresponding to an amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2 is provided with not less than 85% of homology to the amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2.

[0025] The protein of this invention can be produced by the following steps: a DNA of this invention is inserted into a known expression vector to construct a recombinant plasmid; transformed cells are obtained by introducing this recombinant plasmid thereto; the transformed cells are cultured in a suitable medium to allow a protein of the invention to form and accumulate in the culture; and said protein is harvested from the culture.

[0026] Host-vector systems ordinarily used to express exogenous proteins can be employed as cell and expression vectors. For example, mentioned are a combination of a prokaryotic cell such as *E. coli* and an expression vector adapted to the cell and a combination of a eukaryotic cell such as a mammalian cell and an expression vector adapted to the cell. Culture media and culturing conditions may appropriately be selected in accord with the cells to be employed.

[0027] The proteins of this invention may be expressed as fusion proteins with other proteins. Also, the proteins of the invention may be expressed in their full-length, or alternatively, portions thereof may be expressed as partial peptides.

[0028] Cultures, as used herein, are media and cells in the media. Harvesting proteins of this invention from the cultures can be carried out according to known protein purification methods which utilize, among others, the above-mentioned activity of proteins of the invention as an index.

[0029] Antibodies capable of binding to the proteins of this invention can be prepared according to standard methods using the proteins of the invention as antigens; the antibodies may be referred to as "antibody (ies) of this (the) invention." The antibodies of this invention may be monoclonal antibodies or polyclonal antibodies.

[0030] The protein of this invention may be used intact as an antigen; however, it is preferred that the protein be conjugated to keyhole limpet hemocyanine, bovine serum albumin, egg white albumin, etc. and/or be combined with adjuvants for use as the antigen.

[0031] An animal to be immunized such as a mouse, rabbit, guinea pig, or sheep is immunized by administration of the antigen described above via percutaneous, intraperitoneal, or intravenous injections or the like. A polyclonal antibody can, for example, be obtained by collecting serum from the immunized animal.

[0032] A monoclonal antibody can, for example, be obtained in the following manner. After an animal to be immunized such as a mouse, rabbit, guinea pig, or sheep is immunized by administration of the antigen described above via percutaneous, intraperitoneal, or intravenous injections or the like, its spleen or lymph node is extracted. Cells taken from this are fused with myeloma cells, which are preferably derived from the same animal species as that of the immunized animal, to create a hybridoma. A cell strain is selected by repeating screening and cloning from the obtained hybridoma: the strain incessantly produces an antibody specific to the above-mentioned antigen. A monoclonal antibody is produced in a suitable medium by culturing the thus-selected cell strain in the medium; or alternatively, it is produced in an ascite fluid or the like by culturing the strain in vivo, such as mouse abdominal cavities.

[0033] Purification methods for the resulting polyclonal and monoclonal antibodies include salting out with ammonium sulfate, ion-exchange chromatography using a DEAE cellulose column or the like, affinity chromatography using a Protein A column, and immunoabsorption chromatography. The antibodies of this invention can be detected by immunoassays using the proteins of the invention or labeled antibodies.

[0034] The antibodies of this invention may be fragmented ones, so long as they retain antigen-binding sites (Fab). Specifically mentioned as the present fragmented antibody is a fragment containing Fab that can be obtained by digesting the present antibody with a protease such as papain which does not digest the antigen-binding site.

[0035] The antibodies of this invention may be labeled by being bound to labeling substances. The labeling substances are not particularly limited, so long as they can ordinarily be used in labeling of proteins; and mentioned are enzymes, isotopes, fluorescent substances, etc.

[0036] Next, the culturing method of this invention will be explained. The method of the invention is characterized in that it comprises monolayer culture of chondrocytes in the presence of a membrane-permeable cAMP analog in an amount sufficient to cause the chondrocytes to maintain a differentiated state thereof as cartilage.

[0037] Monolayer culture of chondrocytes can be carried out in a manner similar to the conventional monolayer culture of chondrocytes, except that it is to be done in the presence of the membrane-permeable cAMP analog. For example, the media that are used for culture include  $\alpha$ -modified Eagle's medium containing fetal bovine serum, ascorbic acid, an antibiotic, etc. as appropriate.

[0038] The membrane-permeable cAMP analog is an analog of cAMP that has the ability to permeate the membrane without impairing its function as a so-called second messenger of cAMP; preferably, it is dibutyryl cAMP.

[0039] The quantity of the membrane-permeable cAMP analog present in a medium may be such that it is sufficient to cause the chondrocytes to maintain a differentiated state thereof as cartilage. For example, in the case of dibutyryl cAMP, it is preferably 0.3-0.5 mM.

[0040] As used in the present specification, the "differentiated state" means that the chondrocytes possess at least the properties described in the following (1) to (2):

- (1) Exhibit a spherical shape and are abundant in extracellular matrix; and
- (2) Can be stained with toluidine blue satisfactorily.

[0041] The membrane-permeable cAMP analog in an amount sufficient to cause chondrocytes to maintain a differentiated state thereof as cartilage can also induce the differentiation of dedifferentiated chondrocytes.

[0042] Furthermore, this invention provides human chondrocytes that have been cultured by the culturing method of the invention and that possess the properties described in the following (1) to (3):

- (1) Exhibit a spherical shape and are abundant in extracellular matrix;
- (2) Can be stained with toluidine blue satisfactorily; and
- (3) DNA of this invention is expressed therein.

[0043] Since toluidine blue selectively stains sulfated proteoglycan, the chondrocytes of this invention synthesize

the sulfated proteoglycan.

[0044] These chondrocytes express mRNAs for collagen of Type I and Type II as well as those for aggrecan.

#### EXAMPLES

[0045] This invention will be explained by way of examples hereinbelow.

#### (EXAMPLE 1) Culture of Chondrocytes in Differentiated State

[0046] Epiphyseal cartilage of the femur knee joint of a human fetus that was naturally aborted about 25 weeks of pregnancy (obtained from Norman Bethune University of Medical Sciences, Department of Pathology) was obtained. Chondrocytes were isolated from this cartilage according to the same method as that described in Shimomura et al. *Calcif. Tissue Res.* 19, 179-187 (1975), except that the finely cut cartilage was incubated in a  $\alpha$ -modified Eagle's medium ( $\alpha$ -MEM) containing 3 mg/ml collagenase (Type IA, Sigma) for 3h. The cells were seeded at  $1 \times 10^5$  cells per Type I collagen-coated dish and maintained in  $\alpha$ -MEM (10 ml/dish) containing 10% fetal bovine serum, 50  $\mu$ g/ml of ascorbic acid, 32 unit/ml of penicillin, and 40  $\mu$ g/ml of streptomycin. Dibutyl cAMP (dbcAMP) (1 mM) was added to the culture medium when the cells became subconfluent. Cells were cultured over two days either in the presence or in the absence of dbcAMP. Then, while the cells were harvested, the morphological change of cells was examined. After fixing with ethanol, the cells were stained with toluidine blue.

[0047] The chondrocytes cultured in the presence of dbcAMP exhibited a spherical shape and were abundant in extracellular matrix, whereas the chondrocytes cultured in its absence were fibroblast-like and have a spindle shape, and were deficient in extracellular matrix. Fig. 2A shows a photomicrograph of the morphology of cells on the 6th day after addition of dbcAMP (a-no addition of dbcAMP, b-addition of dbcAMP).

[0048] In the staining with toluidine blue that should selectively stain sulfated proteoglycan, the chondrocytes cultured in the presence of dbcAMP were stained satisfactorily, whereas the chondrocytes cultured in its absence were hardly stained. Fig. 2B shows the results of staining with toluidine blue of cells on the 12th day after addition of dbcAMP (a: no addition of dbcAMP, b: addition of dbcAMP).

[0049] Expression of the mRNA for Type I and Type II collagen, and aggrecan, which serve as molecular markers, was investigated by RT-PCR. Expressions of the molecular markers were compared in the presence of dbcAMP as well as in its absence; as a result, it was indicated that the differentiated state was maintained in the presence of dbcAMP.

[0050] Accordingly, it was recognized that the chondrocytes cultured in the presence of dbcAMP maintained a differentiated state thereof as cartilage (namely, the maintenance of a differentiated phenotype).

[0051] When dose-dependence of the above-noted effect by dbcAMP was studied, said effect increased in a dose-dependent manner, reaching its maximum at 0.3-0.5 mM.

[0052] It has been reported that bFGF and TGF- $\beta$  stabilize or stimulate the expression of rabbit or chicken chondrocytes of the differentiated phenotype. Employing bFGF (0.4 ng/ml) and TGF- $\beta$  (3 ng/ml), the chondrocytes were cultured similarly to above, but they did not maintain their differentiated phenotype in human chondrocytes.

#### (EXAMPL 2) Expression of Specific Genes in the Differentiated Chondrocytes

[0053] Total RNA was extracted from the chondrocytes cultured in the presence of dbcAMP and those cultured in the absence of dbcAMP (as described in Example 1) by the guanidiniethiocyanate/cesium trifluoroacetate method. Poly(A)<sup>+</sup>RNA was concentrated using Oligotex-dT30 (Roche). The subtractive hybridization was used to select clones in which mRNA was observed that had been expressed in the differentiated chondrocytes (+dbcAMP) but not in the dedifferentiated chondrocytes (-dbcAMP). Using a PCR select cDNA subtraction kit (Clontech), cDNA synthesized from the mRNA of the differentiated chondrocytes was allowed to hybridize with an excess amount of cDNA of from the mRNA of the dedifferentiated chondrocytes. The cDNA that did not hybridize, namely that was expressed in a differentiated state was amplified by suppression PCR according to the manufacturer's manual. The resulting PCR product was cloned into pGEM-T (Promega), a T tail vector, and the nucleotide sequence determination was carried out on about 120 clones. One clone (pSUB37) was selected for further analysis, and the corresponding protein product was named DEC1.

[0054] NcoI-PstI fragment from pSUB37 was used as a probe to study the expression of DEC1 mRNA in various human fetus tissues by Northern blot analysis. Consequently, DEC1 was expressed in the cartilage, spleen, intestine, and lung; and it was also expressed in the heart, liver, brain, and stomach, although in small amounts. The Northern blot analysis was conducted in the following manner. Total RNA samples (5 or 10  $\mu$ g) were electrophoresed on a 1% agarose gel containing formaldehyde and were transferred to Highbond-N-membranes (Amersham). Total RNA samples of various human fetus tissues were provided by Dr. Li Yu at Norman Bethune University of Medical Sciences, which were intended for the study of tissue distribution. NcoI-PstI fragment from pSUB37 was labeled with [<sup>32</sup>P]dCTP.

and it was used as a hybridization probe. The membranes were washed with 2xSSC containing 0.5% SDS at 65 °C for 30 min. Biomax X-ray films were exposed to the washed membranes using sensitizing films at -70 °C.

[0055] The full-length nucleotide sequence of DEC1 cDNA was determined in the following manner. A full-length cDNA of DEC1 was isolated by RACE (rapid amplification cDNA ends method) using a Marathon cDNA amplification kit (Clontech). Specifically, a double-stranded cDNA was ligated to a Marathon cDNA adapter and subjected to suppression PCR. Reaction was carried out using an adapter primer and a gene-specific primer that had been designed for DEC 1 based on the nucleotide sequence of pSUB37. The amplified cDNA sample was separated on a 4% polyacrylamide gel, DNA of the main band was extracted from the gel, and it was subcloned into pGEM-T. Double-stranded DNA of the subcloned plasmid and a series of synthetic oligonucleotides were used as a sequence-determining template and as a specific primer, respectively. DNA sequence determination was carried out by the Sanger method using either a sequenase 7-deaza-dGTPDNA sequencing kit (Amersham) or an ABI PRISM 310 autosequencer (Perkin-Elmer).

[0056] The nucleotide sequence of DEC1 cDNA thus determined and the amino acid sequence deduced therefrom are set forth in SEQ ID NO: 1. This amino acid sequence alone is also set forth in SEQ ID NO: 2. DEC1 cDNA has an open reading frame of 1236 bp. A 2922 bp length, excluding the poly A region, is well in accord with the size of mRNA (3.1 kb) that was obtained by the Northern blot analysis described above. Since there is a stop codon in the 5'-region which serves as an inframe, the first ATG is regarded as an initiation codon. The sequence around the first ATG coincides with a Kozak consensus sequence (GCCGCCA/GCCATGG). Thus, DEC1 comprises 412 amino acids and its calculated molecular weight is 45.5 kDa.

#### (EXAMPLE 3)

##### (1) Materials and Methods

[0057] Chondrocytes were isolated from the rib growth plate and resting cartilage of ribs of a male Japanese white rabbit (four-weeks old) according to the method as already reported in Shimomura et al. *Calcif. Tissue Res.* 19, 179-187 (1975).

[0058] These cells were seeded at  $5 \times 10^5$  cells per 10 mm plastic culture dish and maintained in 10 ml of  $\alpha$ -MEN supplemented with 10% FBS, 60 mg/ml of kanamycins, 250 ng/ml of amphotericin B, and 50 unit/ml of penicillin G at 37 °C in the air containing 5% CO<sub>2</sub>. After cells reached confluent, the cells were washed with PBS, transferred to a fresh  $\alpha$ -MEN (10 ml) containing no serum, and maintained for 48 h. From 1 to 24 h before the completion of incubation, 1 mM dbcAMP or  $10 \times 10^{-7}$  M human recombinant PTH-(1-84) was added to the medium.

[0059] Human embryonic pulmonary fibroblasts (MRC-5), human cervix uteri epithelial cells (Hela), human hepatoma cells (HepG2) and canine renal epithelial cells (MDCK) were obtained from the Gene Bank of The Institute of Physical and Chemical Research. Cells were cultured in modified Dalbeco Eagle's media (DMEM) supplemented with 10% FBS until they reached confluent. After the cultures had turned confluent, the cells were washed with 10% PBS, transferred to fresh DMEM (10 ml) containing no serum, and maintained for 48 h. From 1 to 24 h before the completion of incubation, 1 mM dbcAMP was added to the media. Then, cells were harvested for RNA preparations.

##### (2) Northern Blot Analysis

[0060] Total RNAs were extracted from the cultured cells by the guanidine thiocyanate/cesium trifluoroacetate method. (Smale G. and Sasse J., *Anal. Biochem.* 203, 352-356 (1992).) The total RNA samples (8-20  $\mu$ g) were electrophoresed on a 1% agarose gel containing formaldehyde and were transferred to NYTRAN membranes (Schleicher & Schuell, Japan). A 1.1 kb NcoI-PstI fragment from pSUB37 was labeled with [<sup>32</sup>P]dCTP, and it was used as a hybridization probe. The membranes were washed with 0.2xSSC containing 0.5% DSD at 55 °C for 30 min. BioMax X-ray films (Eastman Kodak Co., Rochester, NY) were exposed to the washed membranes using sensitizing screens at -70 °C.

##### (3) Results

##### [0061]

(I) In human pulmonary fibroblast cell line MRC5, DEC1 mRNA was induced from 1 to 24 h after addition of dbcAMP (Fig. 3).

(II) In the Hela cells originating in human uterine cancer, DEC1 mRNA was also induced within 1 h after addition of dbcAMP (Fig. 4).

(III) In the rabbit chondrocytes, DEC1 mRNA was induced from 1 to 24 h after addition of PTH (Fig. 5).

(IV) In the rabbit chondrocyte culture system, it was also confirmed that DEC1 mRNA had been induced after addition of dbcAMP (Fig. 6)

(v) In the HepG2 cell strain originating in human hepatic cell line, no change in the level of DEC1 mRNA was observed between 1 to 6 h after addition of dbcAMP.

(VI) In the renal cell strain, the addition of dbcAMP also induced DEC1 mRNA (Fig. 7).

[0062] The above results have indicated that DEC1bHLH transcription factor is involved in the mechanism of action of PTH/PTH-rp in chondrocytes. Furthermore, the DEC1bHLH transcription factor was induced in response to the cAMP within one hour in almost all the mesenchymal and epithelial cells tested. This has suggested that the present transcription factor almost universally participates in the gene expression of the cAMP signal system.

#### INDUSTRIAL APPLICABILITY

[0063] As has been explained above, according to this invention there are provided a gene specifically expressed in differentiated human chondrocytes, a method for culturing human chondrocytes in a differentiated state, and human chondrocytes that have been cultured by the method. These are important not only in the analysis of differentiation and degeneration of cartilage, but also in the development of gene therapy for osteoarthritis and rheumatoid arthritis. Moreover, they are believed to be useful in the treatment of diseases involving other cAMP systems, because cAMP induces DEC1 mRNA in a large number of cells other than the chondrocytes.



## SEQUENCE LISTING

5

SEQ ID NO: 1

LENGTH: 2948

10

TYPE: nucleic acid

STRANDEDNESS: double-stranded

15

TOPOLOGY: linear

MOLECULE TYPE: cDNA

ORIGINAL SOURCE: human

20

CELL TYPE: chondrocyte

FEATURES:

25

NAME/KEY: CDS

LOCATION: 207-1442

SEQUENCE DESCRIPTION:

30

ATTACGAAGT GGACACCGGG CCATGCACGC CCCCAACTGA AGCTGCATCT CAAAGCCGAA 60

35

GATTCCAGCA CCCGAGGGGA TTTCAAAGAG CTCAGACTCA GAGGAACATC TCCGGAGAGA 120

CCCCCGAAGC CCTCTCCAGG GCAGTCCTCA TCCAGACGCT CCGCTAGTCC AGACAGGAGC 180

GCGCAGTGGC CCCGGCTCGC CGCGCC ATG GAG CGG ATC CCC AGC GCG CAA CCA 233

40

Met Glu Arg Ile Pro Ser Ala Gln Pro

1

5

45

CCC CCC GCC TGC CTG CCC AAA GCA CCG GGA CTG GAG CAC GGA GAC CTA 281

Pro Pro Ala Cys Leu Pro Lys Ala Pro Gly Leu Glu His Gly Asp Leu

10

15

20

25

50

CCA GGG ATG TAC CCT GCC CAC ATG TAC CAA GTG TAC AAG TCA AGA CGG 329

Pro Gly Met Tyr Pro Ala His Met Tyr Gln Val Tyr Lys Ser Arg Arg

55

EP 1 020 517 A1

	30	35	40	
5	GGA ATA AAG CGG AGC GAG GAC AGC AAG GAG ACC TAC AAA TTG CCG CAC	377		
	Gly Ile Lys Arg Ser Glu Asp Ser Lys Glu Thr Tyr Lys Leu Pro His			
	45	50	55	
10	CGG CTC ATC GAG AAA AAG AGA CGT GAC CGG ATT AAC GAG TGC ATC GCC	425		
	Arg Leu Ile Glu Lys Lys Arg Arg Asp Arg Ile Asn Glu Cys Ile Ala			
	60	65	70	
15	CAG CTG AAG GAT CTC CTA CCC GAA CAT CTC AAA CTT ACA ACT TTG GGT	473		
	Gln Leu Lys Asp Leu Leu Pro Glu His Leu Lys Leu Thr Thr Leu Gly			
20	75	80	85	
	CAC TTG GAA AAA GCA GTG GTT CTT GAA CTT ACC TTG AAG CAT GTG AAA	521		
25	His Leu Glu Lys Ala Val Val Leu Glu Leu Thr Leu Lys His Val Lys			
	90	95	100	105
	GCA CTA ACA AAC CTA ATT GAT CAG CAG CAG CAG AAA ATC ATT GCC CTG	569		
30	Ala Leu Thr Asn Leu Ile Asp Gln Gln Gln Gln Lys Ile Ile Ala Leu			
	110	115	120	
35	CAG AGT GGT TTA CAA GCT GGT GAG CTG TCA GGG AGA AAT GTC GAA ACA	617		
	Gln Ser Gly Leu Gln Ala Gly Glu Leu Ser Gly Arg Asn Val Glu Thr			
	125	130	135	
40	GGT CAA GAG ATG TTC TGC TCA GGT TTC CAG ACA TGT GCC CGG GAG GTG	665		
	Gly Gln Glu Met Phe Cys Ser Gly Phe Gln Thr Cys Ala Arg Glu Val			
	140	145	150	
45	CTT CAG TAT CTG GCC AAG CAC GAG AAC ACT CGG GAC CTG AAG TCT TCG	713		
	Leu Gln Tyr Leu Ala Lys His Glu Asn Thr Arg Asp Leu Lys Ser Ser			
50	155	160	165	
	CAG CTT GTC ACC CAC CTC CAC CGG GTG GTC TCG GAG CTG CTG CAG GGT	761		

55

EP 1 020 517 A1

	Gln Leu Val Thr His Leu His Arg Val Val Ser Glu Leu Leu Gln Gly	
5	170                      175                      180                      185	
	GGT ACC TCC AGG AAG CCA TCA GAC CCA GCT CCC AAA GTG ATG GAC TTC	809
	Gly Thr Ser Arg Lys Pro Ser Asp Pro Ala Pro Lys Val Met Asp Phe	
10	190                      195                      200	
	AAG GAA AAA CCC AGC TCT CCG GCC AAA GGT TCG GAA GGT CCT GGG AAA	857
15	Lys Glu Lys Pro Ser Ser Pro Ala Lys Gly Ser Glu Gly Pro Gly Lys	
	205                      210                      215	
	AAC TGC GTG CCA GTC ATC CAG CGG ACT TTC GCT CAC TCG AGT GGG GAG	905
20	Asn Cys Val Pro Val Ile Gln Arg Thr Phe Ala His Ser Ser Gly Glu	
	220                      225                      230	
25	CAG AGC GGC AGC GAC ACG GAC ACA GAC AGT GGC TAT GGA GGA GAA TCG	953
	Gln Ser Gly Ser Asp Thr Asp Thr Asp Ser Gly Tyr Gly Gly Glu Ser	
	235                      240                      245	
30	GAG AAG GGC GAC TTG CGC AGT GAG CAG CCG TGC TTC AAA AGT GAC CAC	1001
	Glu Lys Gly Asp Leu Arg Ser Glu Gln Pro Cys Phe Lys Ser Asp His	
35	250                      255                      260                      265	
	GGA CGC AGG TTC ACG ATG GGA GAA AGC ATC GGC GCA ATT AAG CAA GAG	1049
	Gly Arg Arg Phe Thr Met Gly Glu Arg Ile Gly Ala Ile Lys Gln Glu	
40	270                      275                      280	
	TCC GAA GAA CCC CCC ACA AAA AAG AAC CGG ATG CAG CTT TCG GAT GAT	1097
45	Ser Glu Glu Pro Pro Thr Lys Lys Asn Arg Met Gln Leu Ser Asp Asp	
	285                      290                      295	
	GAA GGC CAT TTC ACT AGC AGT GAC CTG ATC AGC TCC CCG TTC CTG GGC	1145
50	Glu Gly His Phe Thr Ser Ser Asp Leu Ile Ser Ser Pro Phe Leu Gly	
	300                      305                      310	

55

EP 1 020 517 A1

CCA CAC CCA CAC CAG CCT CCT TTC TGC CTG CCC TTC TAC CTG ATC CCA 1193  
5 Pro His Pro His Gln Pro Pro Phe Cys Leu Pro Phe Tyr Leu Ile Pro  
315 320 325  
CCT TCA GCG ACT GCC TAC CTG CCC ATG CTG GAG AAG TGC TGG TAT CCC 1241  
10 Pro Ser Ala Thr Ala Tyr Leu Pro Met Leu Glu Lys Cys Trp Tyr Pro  
330 335 340 345  
ACC TCA GTG CCA GTG CTA TAC CCA GGC CTC AAC GCC TCT GCC GCA GCC 1289  
15 Thr Ser Val Pro Val Leu Tyr Pro Gly Leu Asn Ala Ser Ala Ala Ala  
350 355 360  
CTC TCT AGC TTC ATG AAC CCA GAC AAG ATC TCG GCT CCC TTG CTC ATG 1337  
20 Leu Ser Ser Phe Met Asn Pro Asp Lys Ile Ser Ala Pro Leu Leu Met  
365 370 375  
CCC CAG AGA CTC CCT TCT CCC TTG CCA GCT CAT CCG TCC GTC GAC TCT 1385  
25 Pro Gln Arg Leu Pro Ser Pro Leu Pro Ala His Pro Ser Val Asp Ser  
380 385 390  
TCT GTC TTG CTC CAA GCT CTG AAG CCA ATC CCC CCT TTA AAC TTA GAA 1433  
30 Ser Val Leu Leu Gln Ala Leu Lys Pro Ile Pro Pro Leu Asn Leu Glu  
395 400 405  
ACC AAA GAC TAAACTCTCT AGGGGATCCT GCTGCTTTGC TTTCCTTCCT 1482  
40 Thr Lys Asp  
410  
CGCTACTTCC TAAAAAGCAA CAAAAAGTT TTGTGAATG CTGCAAGATT GTTGCAATTGT 1542  
45 GTATACTGAG ATAATCTGAG GCATGGAGAG CAGATTCAGG GTGTGTGTGT GTGTGTGTGT 1602  
GTGTGTGTGT ATGTCCGTGT GCGTGCACAT GTGTGCCTGC GTGTTGGTAT AGGACTTTAA 1662  
50 AGCTCCTTTT GGCATAGGGA AGTCACGAAG GATTGCTTGA CATCAGGAGA CTTGGGGGGG 1722  
ATGTAGCAG ACGTCTGGGC TTTTCCCCAC CCAGAGAATA GCCCCCTTCG ATACACATCA 1782  
55

5 GCTGGATTTT CAAAAGCTTC AAAGTCTTGG TCTGTGAGTC ACTCTTCACT TTGGGAGCTG 1842  
GGTCTGTGGC TTTGATCAGA AGGTACTTTC AAAAGAGGGC TTTCCAGGGC TCAGCTCCCA 1902  
ACCAAGCTGT AGGACCCAC CCTTTTGCCT TTATTGTGCA CGTGACTCAC CAGACGTCGG 1962  
10 GGAGAGAGAG CAGTCAGACC GAGCTTTCTG CTAACATGGG GAGGTAGCAG GCACCTGGCAT 2022  
AGCACGGTAG TGGTTTGGGG AGGTTTCCGC AGGTCTGCTC CCCACCCCTG CCTCGGAAGA 2082  
ATAAAGAGAA TGTAGTCCC TACTCAGGCT TTCGTAGTGA TTAGCTTACT AAGGAACTGA 2142  
15 AAATGGGCCC CTGTACAAG CTGAGCTGCC CCGGAGGGAG GGAGGAGTTC CCTGGGCTTC 2202  
TGGCACCTGT TTCTAGGCCT AACCATTAGT ACTTACTGTG CAGGGAACCA AACCAAGGTC 2262  
TGAGAAATGC GGACACCCCG AGCGAGCACC CCAAAGTGCA CAAAGCTGAG TAAAAAGCTG 2322  
20 CCCCCCTCAA ACAGAACTAG ACTCAGTTT CAATCCATC CTAAACTCC TTTAACCAA 2382  
GCTTAGCTTC TCAAAGGCTT AACCAAGCCT TGGCACCGCC AGATCCTTC TGTAGGCTAA 2442  
25 TTCCTCTGCG CCAACGGCAT ATGGAGTGTC CTTATTGCTA AAAAGGATTC CGTCTCCTTC 2502  
AAAGAAGTTT TATTTTGGT CCAGAGTACT TGTTTCCCG ATGTGTCCAG CCAGCTCCGC 2562  
AGCAGCTTTT CAAGATGCAC TATCCCTGAT TGCTGATCGT GTTTTAACCT TTTCTTTTC 2622  
30 TGTTTTATT TTGGTATTAA GTCGTGCCT TTATTGTAA AGCTGTATA AATATATATT 2682  
ATATAATAT ATTAAGAAG AAAATGTTT AGATGTTTAT TTGTATAATT ACTTGATTCA 2742  
35 CACAGTGAGA AAAAATGAAT GTATTCCTGT TTTGAAGAG AAGAATAATT TTTTTTCTC 2802  
TAGGGAGAGG TACAGTGTTC ATATTTTGA GCCTTCCTGA AGGTGTAAAA TTGTAAATAT 2862  
TTTTATCTAT GAGTAAATGT TAAGTAGTTG TTTTAAATA CTTAATAAAA TAATCTTTT 2922  
40 CCTGTGGAAG AAAAAAAAAA AAAAAA 2948

SEQ ID NO: 2

LENGTH: 412

TYPE: amino acid

TOPOLOGY: linear

MOLECULE TYPE: protein

## SEQUENCE DESCRIPTION:

5

Met Glu Arg Ile Pro Ser Ala Gln Pro Pro Pro Ala Cys Leu Pro Lys

1 5 10 15

10

Ala Pro Gly Leu Glu His Gly Asp Leu Pro Gly Met Tyr Pro Ala His

20 25 30

15

Met Tyr Gln Val Tyr Lys Ser Arg Arg Gly Ile Lys Arg Ser Glu Asp

35 40 45

Ser Lys Glu Thr Tyr Lys Leu Pro His Arg Leu Ile Glu Lys Lys Arg

20

50 55 60

Arg Asp Arg Ile Asn Glu Cys Ile Ala Gln Leu Lys Asp Leu Leu Pro

25

65 70 75 80

Glu His Leu Lys Leu Thr Thr Leu Gly His Leu Glu Lys Ala Val Val

85 90 95

30

Leu Glu Leu Thr Leu Lys His Val Lys Ala Leu Thr Asn Leu Ile Asp

100 105 110

Gln Gln Gln Gln Lys Ile Ile Ala Leu Gln Ser Gly Leu Gln Ala Gly

35

115 120 125

Glu Leu Ser Gly Arg Asn Val Glu Thr Gly Gln Glu Met Phe Cys Ser

40

130 135 140

Gly Phe Gln Thr Cys Ala Arg Glu Val Leu Gln Tyr Leu Ala Lys His

145 150 155 160

45

Glu Asn Thr Arg Asp Leu Lys Ser Ser Gln Leu Val Thr His Leu His

165 170 175

50

Arg Val Val Ser Glu Leu Leu Gln Gly Gly Thr Ser Arg Lys Pro Ser

180 185 190

55

EP 1 020 517 A1

Asp Pro Ala Pro Lys Val Met Asp Phe Lys Glu Lys Pro Ser Ser Pro  
 195 200 205  
 Ala Lys Gly Ser Glu Gly Pro Gly Lys Asn Cys Val Pro Val Ile Gln  
 210 215 220  
 Arg Thr Phe Ala His Ser Ser Gly Glu Gln Ser Gly Ser Asp Thr Asp  
 225 230 235 240  
 Thr Asp Ser Gly Tyr Gly Gly Glu Ser Glu Lys Gly Asp Leu Arg Ser  
 245 250 255  
 Glu Gln Pro Cys Phe Lys Ser Asp His Gly Arg Arg Phe Thr Met Gly  
 260 265 270  
 Glu Arg Ile Gly Ala Ile Lys Gln Glu Ser Glu Glu Pro Pro Thr Lys  
 275 280 285  
 Lys Asn Arg Met Gln Leu Ser Asp Asp Glu Gly His Phe Thr Ser Ser  
 290 295 300  
 Asp Leu Ile Ser Ser Pro Phe Leu Gly Pro His Pro His Gln Pro Pro  
 305 310 315 320  
 Phe Cys Leu Pro Phe Tyr Leu Ile Pro Pro Ser Ala Thr Ala Tyr Leu  
 325 330 335  
 Pro Met Leu Glu Lys Cys Trp Tyr Pro Thr Ser Val Pro Val Leu Tyr  
 340 345 350  
 Pro Gly Leu Asn Ala Ser Ala Ala Ala Leu Ser Ser Phe Met Asn Pro  
 355 360 365  
 Asp Lys Ile Ser Ala Pro Leu Leu Met Pro Gln Arg Leu Pro Ser Pro  
 370 375 380  
 Leu Pro Ala His Pro Ser Val Asp Ser Ser Val Leu Leu Gln Ala Leu  
 385 390 395 400

Lys Pro Ile Pro Pro L u Asn Leu Glu Thr Lys Asp

5

405

410

10

SEQ ID NO: 3

LENGTH: 4

TYPE: amino acid

15

TOPOLOGY: linear

MOLECULE TYPE: peptide

20

SEQUENCE DESCRIPTION:

Trp Arg Pro Trp

25

1

# Claims

30

1. A DNA encoding a protein defined in the following (a) or (b):

(a) a protein comprising the amino acid sequence set forth in SEQ ID NO: 2;

35

(b) a protein comprising an amino acid sequence derived from the amino acid sequence set forth in SEQ ID NO: 2 by deletion or substitution of one or more amino acids therefrom, or by addition of one or more amino acids and capable of binding to nucleotide sequence CANNTG and/or nucleotide sequence CACNAG upon formation of a dimer,

40

wherein the amino acid sequence of a part of said protein corresponding to an amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2 is provided with not less than 85% of homology to the amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2.

2. A DNA represented by the following (c) or (d):

45

(c) a DNA comprising a nucleotide sequence of from nucleotide no. 207 to nucleotide no. 1442 of the nucleotide sequence set forth in SEQ ID NO: 1 or a complementary nucleotide sequence thereto;

(d) a DNA capable of hybridization to the DNA defined in (c) under stringent conditions.

3. A protein represented by the following (e) or (f):

50

(e) a protein comprising the amino acid sequence set forth in SEQ ID NO: 2;

(f) a protein comprising an amino acid sequence derived from the amino acid sequence set forth in SEQ ID NO: 2 by deletion or substitution of one or more amino acids therefrom, or by addition of one or more amino acids thereto and capable of binding to nucleotide sequence CANNTG and/or nucleotide sequence CACNAG upon formation of a dimer,


55

wherein the amino acid sequence of a part of said protein corresponding to an amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2 is provided with not less than 85% of homology to the amino acid sequence of from amino acid no. 51 to amino acid no. 108 in SEQ ID NO: 2.




4. An antibody capable of binding to the protein according to claim 3.
5. A method for culturing human chondrocytes, which comprises monolayer-culturing the chondrocytes in the presence of a membrane-permeable cAMP analog in an amount sufficient to cause the chondrocytes to maintain a differentiated state thereof as cartilage.
6. The method for culturing according to claim 5, wherein the membrane-permeable cAMP analog is dibutyl cAMP.
7. Human chondrocytes that have been cultured by the culturing method according to claim 5 or claim 6, possessing the properties defined in the following (1)-(3):
- (1) exhibiting a spherical shape and being abundant in extracellular matrix;
  - (2) capable of being satisfactorily stained with toluidine blue; and
  - (3) causing the DNA according to claim 1 to be expressed therein.


Fig.1




***Fig.2A***




**Fig.2B**




**Fig.3**




**Fig.4**




**Fig.5**



**Fig.6**





**Fig.7**

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JF98/03106

## A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.<sup>1</sup> C12N15/12, C12P21/02, C12N5/08, C07K14/47, C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.<sup>1</sup> C12N15/12, C12P21/02, C12N5/08, C07K14/47, C07K16/18

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Genbank/EMBL/DBJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
PX	Biochem. Biophys. Res. Commun., 236, (1997), Ming Shen et al., "Molecular characterization of the novel basic helix-loop-helix protein DEC1 expressed in differentiated human embryo chondrocytes" p.294-298	1-7
X	WO, 96-39427, A (Trustees of dartmouth colleg), 12 December, 1996 (12. 12. 96) & EP, 832117, A	1-4

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is

cited to establish the publication date of another citation or other

special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other

means

"P" document published prior to the international filing date but later than

the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be

considered to involve an inventive step when the document is

combined with one or more other such documents, such combination

being obvious to a person skilled in the art

"A" document member of the same patent family

Date of the actual completion of the international search  
28 October, 1998 (28. 10. 98)Date of mailing of the international search report  
10 November, 1998 (10. 11. 98)Name and mailing address of the ISA/  
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)